

Active Surveillance Follow-Up Calendar Low-Intensity Surveillance Plan

You and your doctor have decided on Active Surveillance as a management strategy for your prostate cancer. During Active Surveillance follow-up evaluations are recommended approximately as often as listed below:

Prostate Specific Antigen (blood test)	Obtain every 12 months
Rectal exam	Obtain every 12 months
Biopsy or MRI	Obtain every 36 months

The following Calendar can be filled in to help you remember when return visits are needed and what repeat testing will be obtained at a given visit.

Date of Diagnosis: ______

Date of Confirmatory Test - Start of Surveillance Phase: _____

Surveillance Schedule

Date:

12 months from confirmatory test and AS decision

Recommended evaluations:

- 1) Clinic visit
- 2) PSA blood test
- 3) Rectal exam

Date:

24 months from confirmatory test and AS decision

Recommended evaluations:

- 1) Clinic visit
- 2) PSA blood test
- 3) Rectal exam

Date:

36 months from confirmatory test and AS decision

Recommended evaluations:

- 1) Clinic visit
- 2) PSA blood test
- 3) Rectal exam
- 4) Biopsy OR MRI

Date:

48 months from confirmatory test and AS decision

Recommended evaluations:

- 1) Clinic visit
- 2) PSA blood test
- 3) Rectal exam

Date:

60 months from confirmatory test and AS decision

Recommended evaluations:

- 1) Clinic visit
- 2) PSA blood test
- 3) Rectal exam

Date:

72 months from confirmatory test and AS decision

Recommended evaluations:

- 1) Clinic visit
- 2) PSA blood test
- 3) Rectal exam
- 4) Biopsy OR MRI

^{*}Surveillance may likely go beyond final date shown, at this point an updated calendar should be provided