



## **2021 Clinical Quality Measures for MIPS**

Date modified: January 6, 2021

2021 QCDR Measure IDs	Measure Title	Measure Description	Denominator	Numerator	Denominator Exclusions	High Priority	Measure Type	NQS Domain	Meaningful Measure Area	Care Setting	Inverse Measure	Risk Adjustment
MUSIC4	Prostate Cancer: Active Surveillance/Watchful Waiting for Low Risk Prostate Cancer Patients	Proportion of patients with low-risk prostate cancer receiving active surveillance or watchful waiting	# of low-risk prostate cancer patients 30 or older	# of low-risk prostate cancer patients on active surveillance or watchful waiting	Prostate cancer patients < 30 years of age; patients that have had prior treatment for prostate cancer	No	Process	Effective Clinical Care	Appropriate use of Healthcare	Outpatient Services	No	N/A
MUSIC5	Prostate Cancer: Radical Prostatectomy Cases LOS	Percentage of radical prostatectomy cases with a length of stay > 2 days	# of patients aged 30 or older who underwent a radical prostatectomy as treatment for prostate cancer	# of patients with a length of stay > 2 days after radical prostatectomy	Prostate cancer patients < 30 years of age; patients that have had prior treatment for prostate cancer	Yes - Outcome	Outcome	Effective Clinical Care	Patient-Focused Episode of Care	Hospital Inpatient	Yes	Yes - Proportion of radical prostatectomies with a LOS greater than 2 days
MUSIC10	Prostate Cancer: Confirmation Testing in low risk AS eligible patients	Percentage of low risk patients that are eligible for active surveillance who receive confirmation testing within 6 months of diagnosis	# of patients aged 30 or older with new diagnosis of low and low-intermediate prostate cancer (Gleason 6 or low volume Gleason 3+4)	# of patients that underwent a second biopsy, MRI, or genomics test within 6 months after date of diagnosis (positive)	Prostate cancer patients < 30 years of age; Patients that have had prior treatment for prostate cancer; Patients on watchful waiting	No	Process	Effective Clinical Care	Appropriate use of Healthcare	Outpatient Services	No	N/A
MUSIC11	Prostate Cancer: Follow-Up Testing for patients on active surveillance for at least 30 months	Percentage of patients on active surveillance that have ≥ 2 tumor burden reassessments and 3 PSA tests in first 30 months since diagnosis	# of patients aged 30 or older with new diagnosis of low and low-intermediate prostate cancer (Gleason 6 or low volume Gleason 3+4)	# of patients on active surveillance that have ≥ 2 tumor burden reassessments and 3 PSA tests in first 30 months since diagnosis	Prostate cancer patients < 30 years of age; Patients that have had prior treatment for prostate cancer	No	Process	Effective Clinical Care	Appropriate use of Healthcare	Outpatient Services	No	N/A
MUSIC12	Kidney Stones: ED visit within 30 days of ureteroscopy	Percentage of patients who underwent ureteroscopy and experienced an unplanned ED visit within 30 days of the procedure	# of patients that undergo ureteroscopy	# of patients with an ED visit within 30 days of ureteroscopy	Patient under 18 at time of surgery; Bilateral surgery; No stones found during URS; Staged surgery for which the primary procedure was not done locally; Other simultaneous procedures (e.g. ureteral biopsy); Nephrostomy tube in place at the time of surgery. Further, due to the significant volume of cases, MUSIC executes a sampling methodology for case entry. MUSIC practices that perform 100-200 cases per year exclude patients born in Mar, Jun, Sep, and Dec and sites with >200 cases per year exclude patients born in Feb, Apr, Jun, Aug, Oct, and Dec	Yes - Outcome	Outcome	Patient Safety	Admissions and Readmissions to Hospitals	Hospital	Yes	Yes - Proportion of ureteroscopies with an ED visit within 60 days of the procedure
MUSIC15	Kidney Stones: SWL in patients with largest renal stone > 2 cm or lower pole stone > 1 cm	Percentage of patients who underwent shockwave lithotripsy with a largest renal stone > 2 cm or with a lower pole stone > 1 cm	# of patients with a renal stone > 2 cm or with a lower pole stone > 1 cm	# of patients with a renal stone > 2 cm or with a lower pole stone > 1 cm and undergo shockwave lithotripsy	Patient under 18 at time of surgery; Bilateral surgery; Staged surgery for which the primary procedure was not done locally; Other simultaneous procedures (e.g. ureteral biopsy); Nephrostomy tube in place at the time of surgery. Further, due to the significant volume of cases, MUSIC executes a sampling methodology for case entry. MUSIC practices that perform 100-200 cases per year exclude patients born in Mar, Jun, Sep, and Dec and sites with >200 cases per year exclude patients born in Feb, Apr, Jun, Aug, Oct, and Dec	No	Process	Effective Clinical Care	Appropriate use of Healthcare	Hospital	Yes	N/A
MUSIC17	Kidney Stones: Opioid utilization after ureteroscopy and shockwave lithotripsy	Percentage of patients who underwent ureteroscopy or shockwave lithotripsy and are discharged on NSAIDs, Acetaminophen, or "Other" and who were not prescribed opioids for pain control	# of patients that undergo ureteroscopy or shockwave lithotripsy	# of patients that were discharged on NSAIDs, Acetaminophen, or "Other" and who were not prescribed opioids for pain control after ureteroscopy or shockwave lithotripsy	Patient under 18 at time of surgery; Bilateral surgery; No stones found during URS; Staged surgery for which the primary procedure was not done locally; Other simultaneous procedures (e.g. ureteral biopsy); Nephrostomy tube in place at the time of surgery; Patients with "unknown" pain medication prescription at discharge. Further, due to the significant volume of cases, MUSIC executes a sampling methodology for case entry. MUSIC practices that perform 100-200 cases per year exclude patients born in Mar, Jun, Sep, and Dec and sites with >200 cases per year exclude patients born in Feb, Apr, Jun, Aug, Oct, and Dec	Yes	Process	Patient Safety	Prevention and Treatment of Opioid and Substance Use Disorders	Hospital	No	N/A
MUSIC18	Kidney Stones: Alpha-blockers at discharge for patients undergoing ureteroscopy or shockwave lithotripsy	Percentage of patients who underwent ureteroscopy or shockwave lithotripsy and received alpha-blockers at discharge	# of patients that undergo ureteroscopy and shockwave lithotripsy	# of patients who undergo ureteroscopy or shockwave lithotripsy and receive alpha blockers at discharge	Patient under 18 at time of surgery; Bilateral surgery; No stones found during URS; Staged surgery for which the primary procedure was not done locally; Other simultaneous procedures (e.g. ureteral biopsy); Nephrostomy tube in place at the time of surgery. Further, due to the significant volume of cases, MUSIC executes a sampling methodology for case entry. MUSIC practices that perform 100-200 cases per year exclude patients born in Mar, Jun, Sep, and Dec and sites with >200 cases per year exclude patients born in Feb, Apr, Jun, Aug, Oct, and Dec	No	Process	Effective Clinical Care	Medication Management	Hospital	No	N/A
MUSIC19	Kidney Stones: Readmission within 30 days of ureteroscopy	Percentage of patients who underwent ureteroscopy and experienced a readmission within 30 days of the procedure	# of patients that undergo ureteroscopy	# of patients with readmission within 30 days of ureteroscopy	Patient under 18 at time of surgery; Bilateral surgery; No stones found during URS; Staged surgery for which the primary procedure was not done locally; Other simultaneous procedures (e.g. ureteral biopsy); Nephrostomy tube in place at the time of surgery; Patients with "unknown" pain medication prescription at discharge. Further, due to the significant volume of cases, MUSIC executes a sampling methodology for case entry. MUSIC practices that perform 100-200 cases per year exclude patients born in Mar, Jun, Sep, and Dec and sites with >200 cases per year exclude patients born in Feb, Apr, Jun, Aug, Oct, and Dec	Yes	Outcome	Patient Safety	Admissions and Readmissions to Hospitals	Hospital	Yes	N/A
MUSIC20	Prostate Cancer: Complications within 30 days of radical prostatectomy	Percentage of radical prostatectomy patients who do not have any deviations from an uncomplicated recovery pathway within 30 days of the procedure	# of patients that undergo radical prostatectomy	# of patients with zero deviations within 30 days of radical prostatectomy	Patients < 30 years of age; patients that have had prior treatment for prostate cancer; patients that are not being screened or treated for prostate cancer	Yes - Outcome	Outcome	Effective Clinical Care	Admissions and Readmissions to Hospitals	Hospital Inpatient	No	Yes - Proportion of patients with a deviation from an "uncomplicated recovery pathway" or non-routine
MUSIC21	Prostate Cancer: Opioid utilization after radical prostatectomy	Percentage of patients who underwent radical prostatectomy and are discharged with ≤ 6 opioid pain pills (5mg oxycodone or equivalent) and do not get a prescription for opioids within 30 days of	# of opioid-naïve patients that undergo radical prostatectomy and complete the baseline and 1 month PRO survey	# of patients that were discharged with ≤ 6 opioid pain pills (5mg oxycodone or equivalent) for pain control and do not get a prescription for opioids within 30 days of	Patients < 30 years of age; patients that have had prior treatment for prostate cancer; patients that are not being screened or treated for prostate cancer; patients reporting as opioid users prior to surgery	Yes - Opioid-related measure	Process	Community/Population Health	Medication Management	Hospital Inpatient	No	N/A
MUSIC22	Prostate Cancer: Urinary continence at 12 months post-radical prostatectomy	Percentage of radical prostatectomy patients that are socially continent (0 - 1 pads per day) at 12 months after surgery	# of patients with good baseline function (0-1 pads per day) that undergo radical prostatectomy and complete the 12 month PRO survey	# of radical prostatectomy patients that are socially continent (0 - 1 pads per day) at 12 months after surgery	Patients < 30 years of age; patients that have had prior treatment for prostate cancer; patients that are not being screened or treated for prostate cancer	Yes - Outcome	Patient Reported Outcome (PRO)	Effective Clinical Care	Functional Outcomes	Hospital Inpatient	No	Yes - Percentage of radical prostatectomy patients that are socially continent (0 - 1 pads per day) at 12 months

All proportional measures; No numerator exclusions or denominator exceptions for MUSIC QCDR measures; All of the MUSIC QCDR approved measures have one performance rate per metric

2021 QCDR Measure IDs	Measure Title	Measure Description	Denominator	Numerator	Denominator Exclusions	High Priority	Measure Type	NQS Domain	Meaningful Measure Area Rationale		Inverse Measure	Risk Adjustment
MUSIC23	Renal Mass: Documentation of the RENAL score for patients with small renal mass diagnoses	Percentage of patients diagnosed with a small renal mass ( $\leq 7$ cm) and have their RENAL score documented in the medical record by the attending	# of patients with a small renal mass diagnosis	# of patients with a RENAL score documented in the medical record	Patients < 18 years of age; Patients with renal mass > 7 cm	Yes - Care Coordination	Process	Communication and Care Coordination	Appropriate use of Healthcare	Outpatient Services	No	N/A
MUSIC24	Kidney Stones: Post-ureteroscopy and shockwave lithotripsy imaging for any stones	Percentage of patients who underwent imaging within 60 days after ureteroscopy or shockwave lithotripsy	# of patients that undergo ureteroscopy or shockwave lithotripsy	# of patients that undergo imaging (Ultrasound and/or CT scan) for ureteral stones within 60 days after ureteroscopy or imaging (Ultrasound, CT scan, and/or Abdominal X-ray) within 60 days after shockwave lithotripsy	Patients under 18 at time of surgery; Bilateral surgery; Staged surgery for which the primary procedure was not done locally; Other simultaneous procedures (e.g. ureteral biopsy); Nephrostomy tube in place at the time of surgery; Patients with "unknown" pain medication prescription at discharge. Further, due to the significant volume of cases, MUSIC executes a sampling methodology for case entry. MUSIC practices that perform 100-200 cases per year exclude patients born in Mar, Jun, Sep, and Dec and sites with >200 cases per year exclude patients born in Feb, Apr, Jun, Aug, Oct, and Dec	No	Process	Effective Clinical Care	Appropriate use of Healthcare	Outpatient Services	No	N/A
MUSIC25	Renal Mass: ED visit or Readmission within 30 days of radical nephrectomy	Percentage of patients with a small renal mass ( $\leq 7$ cm) who underwent a radical nephrectomy and experienced an ED visit or readmission within 30 days	# of patients that undergo radical nephrectomy	# of patients with an ED visit or readmission within 30 days of radical nephrectomy	Patients < 18 years of age; Patients with renal mass > 7 cm	Yes - Outcome	Outcome	Patient Safety	Admissions and Readmissions to Hospitals	Hospital Inpatient	Yes	N/A
MUSIC26	Renal Mass: ED Visit or Readmission within 30 days of partial nephrectomy	Percentage of patients with a small renal mass ( $\leq 7$ cm) who underwent a partial nephrectomy and experienced an ED visit or readmission within 30 days	# of patients that undergo partial nephrectomy	# of patients with an ED visit or readmission within 30 days of partial nephrectomy	Patients < 18 years of age; Patients with renal mass > 7 cm	Yes - Outcome	Outcome	Patient Safety	Admissions and Readmissions to Hospitals	Hospital Inpatient	Yes	N/A
104	Prostate Cancer: Combination Androgen Deprivation Therapy for High Risk or Very High Risk Prostate Cancer	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at high or very high risk of recurrence receiving external beam radiotherapy to the prostate who were prescribed androgen deprivation therapy in combination with external beam radiotherapy to the prostate	Patients with a diagnosis of prostate cancer at high or very high risk of recurrence receiving external beam radiotherapy to the prostat	Patients who were prescribed androgen deprivation therapy in combination with external beam radiotherapy to the prostate	Patients < 30 years of age; patients that; Diagnosis for metastatic cancer Denominator Exception(s): Documentation of medical reason(s) for not prescribing/administering androgen deprivation therapy in combination with external beam radiotherapy to the prostate (e.g., salvage therapy) Documentation of patient reason(s) for not prescribing/administering androgen deprivation therapy in combination with external beam radiotherapy to the prostate	No	Process	Effective Clinical Care	Management of Chronic Conditions	Outpatient Services	No	No
130	Documentation of Current Medications in the Medical Record	Percentage of visits for patients aged 18 years and older for which the eligible professional attests to documenting a list of current medications using all immediate resources available on the date of the encounter.	All visits occurring during the 12 month measurement period	Attestation of documenting, updating or reviewing a patient's current medications using all immediate resources available on the date of encounter.	Patients < 30 years of age; patients that had prior treatment for prostate cancer; Eligible clinician attests to documenting in the medical record the patient is not eligible for a current list of medications being obtained, updated, or reviewed by the eligible clinician	Yes - Patient Safety	Process	Patient Safety	Medication Management	Outpatient Services	No	No
250	Radical Prostatectomy Pathology Reporting	Percentage of radical prostatectomy pathology reports that include the pT category, the pN category, the Gleason score and a statement about margin status	All radical prostatectomy surgical pathology examinations performed during the measurement period for prostate cancer patients	Radical prostatectomy pathology reports that include the pT category, the pN category, Gleason score and a statement about margin status	Patients < 30 years of age; patients that have had prior treatment for prostate cancer; Specimen site other than anatomic location of prostate Denominator Exception(s): Documentation of medical reason(s) for not including pT category, pN category, Gleason score and statement about margin status in the pathology report (e.g., specimen originated from other malignant neoplasms, transurethral resections of the prostate (TURP), or secondary site prostatic carcinomas)	No	Process	Effective Clinical Care	Transfer of Health Information and Interoperability	Hospital	No	No
265	Biopsy Follow-Up	Percentage of new patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician	All new patients undergoing prostate biopsy	Patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and the patient by the provider and/or office and medical team. There must also be acknowledgement and/or documentation of the communication in a biopsy tracking log and document in the patient's	Patients < 30 years of age; patients that have had prior treatment for prostate cancer; patients that are not being screened or treated for prostate cancer; Clinician documented reason that patient's biopsy results were not reviewed, [e.g., patient asks that biopsy results not be communicated to the primary care/referring physician, patient does not have a primary care/referring physician or is a self-referred patient]	Yes - Care Coordination	Process	Communication and Care Coordination	Transfer of Health Information and Interoperability	Outpatient Services	No	No

All proportional measures; No numerator exclusions or denominator exceptions for MUSIC QCDR measures; All of the MUSIC QCDR approved measures have one performance rate per metric