



## **2017 Clinical Quality Measures for MIPS**

Date Modified: June 21, 2017

Measure ID	Measure Title	Measure Description	NQS Domain	Numerator	Denominator	Denominator Exclusions	Denominator Exceptions
MUSIC1	Prostate Biopsy Antibiotic Compliance	Compliance with AUA best practices for antibiotic prophylaxis for transrectal ultrasound-guided (TRUS) biopsy	Patient Safety	# of biopsies where a compliant antibiotic regimen was given	# of prostate biopsies for patients aged 30 or older	Non-TRUS biopsies; biopsies performed in patients < 30 years of age; biopsies performed in patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO and that which have a local biopsy are included, as appropriate, regardless if had have had prior treatment for Pca
MUSIC2	Prostate Biopsy: Unplanned Hospital Admission within 30 Days of TRUS Biopsy	Percentage of biopsies where patients aged 30 years and older had an unplanned hospital admission within 30 days of biopsy	Effective Clinical Care	# of biopsies where a hospital admission resulted within 30 days of the procedure	# of prostate biopsies for patients aged 30 or older	Biopsies performed in patients < 30 years of age; biopsies performed in patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO and that which have a local biopsy are included, as appropriate, regardless if had have had prior treatment for Pca
MUSIC3	Prostate Cancer: Avoidance of Overuse of CT Scan for Staging Low Risk Prostate Cancer Patients	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a CT scan performed at any time since diagnosis of prostate cancer	Efficiency and Cost Reduction	# of low risk prostate cancer patients receiving brachytherapy, EBRT, RP, or cryotherapy who did not have a CT scan since diagnosis of prostate cancer	# of low-risk prostate cancer patients 30 years or older receiving brachytherapy, EBRT, RP, or cryotherapy	Prostate cancer patients < 30 years of age; prostate cancer patients receiving an alternate form of treatment from those specified; patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO are included in the registry, as appropriate, regardless if had have had prior treatment for Pca
MUSIC4	Prostate Cancer: Active Surveillance/Watchful Waiting for Low Risk Prostate Cancer Patients	Proportion of patients with low-risk prostate cancer receiving active surveillance or watchful waiting	Effective Clinical Care	# of low-risk prostate cancer patients on active surveillance or watchful waiting	# of low-risk prostate cancer patients 30 or older	Prostate cancer patients < 30 years of age; patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO are included in the registry, as appropriate, regardless if had have had prior treatment for Pca
MUSIC5	Prostate Cancer: Radical Prostatectomy Cases LOS	Percentage of radical prostatectomy cases with a length of stay > 2 days	Effective Clinical Care	# of patients with a length of stay > 2 days after radical prostatectomy	# of patients aged 30 or older who underwent a radical prostatectomy as treatment for prostate cancer	Prostate cancer patients < 30 years of age; patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO are included in the registry, as appropriate, regardless if had have had prior treatment for Pca
MUSIC6	Prostate Cancer: Unplanned Hospital Readmission After Radical Prostatectomy	Percentage of patients aged 30 years and older who had an unplanned hospital readmission within 30 days of principal procedure	Communication and Care Coordination	# of radical prostatectomies where a hospital admission resulted within 30 days of the procedure	# of radical prostatectomies for patients aged 30 or older	Prostatectomies performed in patients < 30 years of age; patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO are included in the registry, as appropriate, regardless if had have had prior treatment for Pca
MUSIC9	Prostate Biopsy: Repeat Biopsy for Patients with Atypical Small Acinar Proliferation (ASAP)	Proportion of patients undergoing a repeat prostate biopsy within 12 months of their initial biopsy in the registry as a result of a finding of atypical small acinar proliferation (ASAP) as per the NCCN guidelines	Effective Clinical Care	# of patients undergoing a repeat prostate biopsy as a result of a finding of ASAP on their initial biopsy	# of patients aged 30 or older that had a finding of ASAP on their initial prostate biopsy	Patients < 30 years of age; patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO and that which have a local biopsy are included, as appropriate, regardless if had have had prior treatment for Pca
MUSIC10	Prostate Cancer: Confirmation Testing in low risk AS eligible patients	Percentage of low risk patients that are eligible for active surveillance who receive confirmation testing within 6 months of diagnosis	Effective Clinical Care	# of patients that underwent a second biopsy, MRI, or genomics test within 6 months after date of diagnosis (positive biopsy date)	# of patients aged 30 or older with new diagnosis of low and low-intermediate prostate cancer (Gleason 6 or low volume Gleason 3+4)	Prostate cancer patients < 30 years of age; Patients that have had prior treatment for prostate cancer; Patients on watchful waiting	N/A
MUSIC11	Prostate Cancer: Follow-Up Testing for patients on active surveillance for at least 30 months	Percentage of patients on active surveillance that have ≥ 2 tumor burden reassessments and 3 PSA tests in first 30 months since diagnosis	Effective Clinical Care	# of patients on active surveillance that have ≥ 2 tumor burden reassessments and 3 PSA tests in first 30 months since diagnosis	# of patients aged 30 or older with new diagnosis of low and low-intermediate prostate cancer (Gleason 6 or low volume Gleason 3+4)	Prostate cancer patients < 30 years of age; Patients that have had prior treatment for prostate cancer	N/A

102	Prostate Cancer: Avoidance of Overuse of Bone Scan for Staging Low Risk Prostate Cancer Patients	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a bone scan performed at any time since diagnosis of prostate cancer	Efficiency and Cost Reduction	<i>MIPS Quality Measure – See Measure Title/Description</i>	<i>MIPS Quality Measure – See Measure Title/Description</i>	Patients < 30 years of age; patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO are included in the registry, as appropriate, regardless if had have had prior treatment for Pca
130	Documentation of Current Medications in the Medical Record	Percentage of visits for patients aged 18 years and older for which the eligible professional attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration.	Patient Safety	<i>MIPS Quality Measure – See Measure Title/Description</i>	<i>MIPS Quality Measure – See Measure Title/Description</i>	Patients < 30 years of age; patients that have had prior treatment for prostate cancer; patients that are not being screened or treated for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO are included in the registry, as appropriate, regardless if had have had prior treatment for Pca
250	Radical Prostatectomy Pathology Reporting	Percentage of radical prostatectomy pathology reports that include the pT category, the pN category, the Gleason score and a statement about margin status	Effective Clinical Care	<i>MIPS Quality Measure – See Measure Title/Description</i>	<i>MIPS Quality Measure – See Measure Title/Description</i>	Patients < 30 years of age; patients that have had prior treatment for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO are included in the registry, as appropriate, regardless if had have had prior treatment for Pca
265	Biopsy Follow-Up	Percentage of new patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician	Communication and Care Coordination	<i>MIPS Quality Measure – See Measure Title/Description</i>	<i>MIPS Quality Measure – See Measure Title/Description</i>	Patients < 30 years of age; patients that have had prior treatment for prostate cancer; patients that are not being screened or treated for prostate cancer	Prostatectomy patients enrolled in MUSIC PRO and that which have a local biopsy are included , as appropriate, regardless if had have had prior treatment for Pca